

Quantum phases of a Feshbach-resonant atomic Bose gas in one dimension

Yu-Wen Lee

Physics Department, Tunghai University, Taichung, Taiwan, R.O.C.*

Yu-Li Lee

Physics Department, National Changhua University of Education, Changhua, Taiwan, R.O.C.†

We study an atomic Bose gas with an s -wave Feshbach resonance in a one-dimensional optical lattice, with the densities of atoms and molecules incommensurate with the lattice. At zero temperature, most of the parameter region is occupied by a phase in which the superfluid fluctuations of atoms and molecules are the predominant ones, due to the phase fluctuations of atoms and molecules being locked by a Josephson coupling between them. When the density difference between atoms and molecules is commensurate with the lattice, two additional phases may exist: the two component Luttinger liquid where both the atomic and molecular sectors are gapless, and the inter-channel charge density wave where the relative density fluctuations between atoms and molecules are frozen at low energy.

PACS numbers: 03.75.Mn, 03.75.Lm, 67.60.-g

Introduction Trapped dilute cold atomic gases are one of the most exciting fields in condensed matter physics.[1] An important recent development in this area is the application of Feshbach resonances. The energy difference between the molecular state and the two-atom continuum, known as the detuning ν , can be experimentally tuned by means of a magnetic field. Therefore, by sweeping the magnetic field from positive to negative detuning through the Feshbach resonance, it is actually possible to form molecules in the atomic gas.[2]

Depending on the quantum statistics of atoms, the low temperature properties of a dilute atomic gas with an s -wave Feshbach resonance will be much different. In the case of fermionic atoms, a crossover from a superfluid (SF) of the Bardeen-Cooper-Schrieffer type to the Bose-Einstein condensation (BEC) is predicted.[3] On the other hand, a quantum phase transition can occur for bosonic atoms by changing the value of magnetic field detuning.[4] There, two thermodynamically distinct phases exist at zero temperature: the “atomic superfluid” (ASF) phase with both atomic BEC and molecular BEC and the “molecular superfluid” (MSF) with molecular BEC only.

In the present letter, we study the possible quantum phases for an atomic Bose gas with an s -wave Feshbach resonance in a one-dimensional optical lattice. It can be described by the Hamiltonian

$$\begin{aligned} H = & - \sum_i \left(t_m b_{i+1}^\dagger b_i + t_a a_{i+1}^\dagger a_i + \mathcal{H}.c. \right) \\ & - \sum_i [\mu n_{a,i} + (2\mu - \nu) n_{b,i}] \\ & + \sum_i \frac{U_a}{2} n_{a,i}(n_{a,i} - 1) + \frac{U_m}{2} n_{b,i}(n_{b,i} - 1) \\ & + \sum_i U \left(b_i^\dagger a_i a_i + \mathcal{H}.c. \right) + U_{ab} \sum_i n_{a,i} n_{b,i}. \quad (1) \end{aligned}$$

Here $n_{a(b),i}$ is the atom (molecule) density operator and μ is the chemical potential. This system can also be considered a kind of binary mixtures with two types of bosons. However, a main distinction between the present case and the binary mixture studied previously[5, 6] is that a Josephson coupling between atoms and molecules, the U term in Eq. (1), is absent in the latter. As we shall see, this will change the global phase diagram dramatically.

Our main results can be summarized in Fig. 1. For on-site interactions only, i.e. the Hamiltonian given by Eq. (1), the Josephson coupling dominates the low-energy physics in most of the parameter region, and the resulting phase exhibits strong superfluid fluctuations in both atomic and molecular sectors. This is because the phase fluctuations of both sectors are locked by the Josephson coupling. This phase is, in fact, a one-dimensional ($1d$) analog of the ASF phase in three dimensions ($3d$). (But we do not have real condensates here.) Therefore, it will be referred to as $1d$ ASF. The $1d$ ASF will collapse for sufficiently strong attractions between atoms and molecules ($U_{ab} < 0$). When the density difference between atoms and molecules is close to some rational number, depending on the values of parameters as




FIG. 1: Schematic phase diagram of a Feshbach-resonant atomic Bose gas in a one-dimensional optical lattice: (a) on-site interactions only and (b) Tonks limit with nearest-neighbor repulsions between atoms (molecules) V_a (V_m). δ denotes the density difference between atoms and molecules. a_0 is the lattice spacing.

shown in Fig. 1 (b), two additional phases may emerge by including nearest-neighbor repulsions between atoms (molecules): the two component Luttinger liquid (2-LL) where both the atomic and molecular sectors are gapless, and the inter-channel charge density wave (I-CDW) where the relative density fluctuations between atoms and molecules are frozen at low energy.[7]

The continuum theory We now outline below the derivation of our results. For simplicity, we shall consider the case where both the densities of atoms and molecules, $\rho_a = \langle n_{a,i} \rangle / a_0$ and $\rho_b = \langle n_{b,i} \rangle / a_0$, respectively, are incommensurate with the lattice. (Here a_0 is the lattice spacing. ρ_a and ρ_b satisfy the constraint: $\rho_a + 2\rho_b = \rho_0$ where ρ_0 is the density of bare atoms.) That is, we do not consider the possibility of the SF-Mott insulator transition. Then, in terms of the “bosonization” formula[8]: $a_i \sim e^{-i\sqrt{\pi}\theta_a(x)} \sum_{n=-\infty}^{\infty} e^{i2\pi n\rho_a x} e^{i\sqrt{4\pi}n\phi_a(x)}$ and $b_i \sim e^{-i\sqrt{\pi}\theta_m(x)} \sum_{n=-\infty}^{\infty} e^{i2\pi n\rho_b x} e^{i\sqrt{4\pi}n\phi_m(x)}$, the low-energy physics of H [Eq. (1)] can be described by the following effective Hamiltonian:

$$\begin{aligned} H_{eff} = & \frac{v_a}{2} \int dx \left[K_a (\partial_x \theta_a)^2 + \frac{1}{K_a} (\partial_x \phi_a)^2 \right] \\ & + \frac{v_m}{2} \int dx \left[K_m (\partial_x \theta_m)^2 + \frac{1}{K_m} (\partial_x \phi_m)^2 \right] \\ & + g_1 \int dx \cos [\sqrt{\pi}(\theta_m - 2\theta_a)] \\ & + g_2 \int dx \partial_x \phi_a \partial_x \phi_m \\ & + g_3 \int dx \cos [\sqrt{4\pi}(\phi_a - \phi_m) + 2\pi\delta x], \quad (2) \end{aligned}$$

where $\delta = \rho_a - \rho_b$. $v_{a/m}$ and $K_{a/m}$ are sound velocities and Luttinger liquid (LL) parameters, respectively. In Eq (2), only those terms which may become the most relevant in the renormalization group (RG) sense are retained. The values of $v_{a/m}$, $K_{a/m}$, g_1 , g_2 , and g_3 depend on the short-distance physics. In general, they must be extracted from numerics or experiments. $K_{a/m} \gg 1$ in the weak coupling regime. On the other hand, $K_{a/m} = 1$ in the Tonks limit, i.e. $U_{a(m)}/t_{a(m)} \rightarrow +\infty$. Therefore, for on-site interactions only, $1 \leq K_{a/m} < +\infty$. [8] The value of K_a (K_m) can be further decreased by including nearest-neighbor repulsions between atoms (molecules).

The g_3 term is a Umklapp process. It can be neglected when δ is not close to zero (incommensurate filling). On the other hand, when δ is close to zero (commensurate filling), the g_3 term can affect low-energy physics and one may no longer neglect it in Eq. (2). When $a_0|\delta|$ is close to some rational number k/l where $k = 1, 2, \dots$, $l = 2, 3, \dots$, and k and l are co-prime with one another, other Umklapp processes must be taken into account, and one must include the following term in Eq. (2):

$$\tilde{g}_3 \int dx \cos [\sqrt{4\pi}l(\phi_a - \phi_m) + 2\pi l\delta x]. \quad (3)$$

We shall focus on Eq. (2) in the following and discuss the effects of the $\tilde{g}_3$ term later.

Incommensurate filling When δ is incommensurate with the lattice, one may neglect the g_3 term in Eq. (2). To analyze the effects of the g_1 and g_2 terms on low-energy physics, we resort to the RG method. A perturbative calculation up to the one-loop order already shows that the g_1 and g_2 terms alone do not form a closed operator algebra in the sense of operator product expansion (OPE). One must include the term $\partial_x \theta_a \partial_x \theta_m$ in H_{eff} [Eq. (2)]. To simplify the analysis, we consider the case $v_a = v_m = v_0$. (The effects of velocity anisotropy will be discussed later.) Further, we rescale g_1 by $a_0^2 g_1 \rightarrow g_1$. The one-loop RG equations can be obtained by setting $\lambda_3(l) = 0$ in Eqs. (10) – (15). By solving these scaling equations, one may obtain the following results: Within the weak-coupling region, i.e. $\pi|g_1|/(2v_0), |g_2|/(2v_0) \ll 1$, the g_1 term is relevant in the regime $D_1 < 2 + \frac{\sqrt{2\pi}|g_1|}{v_0}$, while it becomes irrelevant for $D_1 > 2 + \frac{\sqrt{2\pi}|g_1|}{v_0}$, where D_1 is the scaling dimension of the g_1 term, defined by

$$D_1 = \frac{1}{K_a} + \frac{1}{4K_m}. \quad (4)$$

Therefore, there are two zero-temperature phases. It turns out that the phase transition between these two phases is of the KT type. We note that for on-site interactions only, i.e. $1 \leq K_{a/m} < +\infty$, the g_1 term is always relevant from Eq. (4).

In the regime where the g_1 term is relevant, it is convenient to define new bosonic fields: $\theta_{\pm} \equiv \theta_a \pm \frac{1}{2}\theta_m$ and $\phi_{\pm} \equiv \frac{1}{2}\phi_a \pm \phi_m$. Then, the value of $\langle \theta_{-} \rangle$ is pinned and a gap $\Delta_0 \sim [\pi|g_1|/(2v_0)]^{2-D_1}$ is opened for the θ_{-} sector. That is, the phase fluctuations of atoms and molecules are locked by the Josephson coupling. By integrating out the gapped sector, the low-energy effective Hamiltonian describing the gapless (θ_{+}) sector takes the form of LLs:

$$H_{+} = \frac{v}{2} \int dx \left[K (\partial_x \theta_{+})^2 + \frac{1}{K} (\partial_x \phi_{+})^2 \right]. \quad (5)$$

For $\pi|g_1|/(2v_0), |g_2|/(2v_0) \ll 1$, v and K can be related to the short distance variables with the help of the one-loop RG equations, yielding

$$\begin{aligned} \frac{v}{v_0} &= \sqrt{\left(\frac{K_a^*}{4} + K_m^* + 2\lambda_4^* \right) \left(\frac{1}{K_a^*} + \frac{1}{4K_m^*} + 2\lambda_2^* \right)}, \\ K &= \sqrt{\frac{K_a^*/4 + K_m^* + 2\lambda_4^*}{1/K_a^* + 1/(4K_m^*) + 2\lambda_2^*}}, \quad (6) \end{aligned}$$

where $K_a^* = K_a + \frac{1}{2-D_1}$, $K_m^* = K_m + \frac{1}{4(2-D_1)}$, $\lambda_2^* = \frac{g_2}{2v_0} + \frac{1}{2K_a K_m (2-D_1)}$, and $\lambda_4^* = -\frac{1}{2(2-D_1)}$. Equation (6) indicates that the 1d ASF will become unstable provided that the inequality is satisfied: $\frac{1}{K_a^*} + \frac{1}{4K_m^*} < -2\lambda_2^*$. This




FIG. 2: Phase diagram for $\delta = 0$: (a) suggested by tree-level scaling analysis and (b) predicted by the one-loop RG equations. $D_1 < 2$ and $D_2 > 2$ in Region I, $D_1 > 2$ and $D_2 < 2$ in Region II, $D_1, D_2 > 2$ in Region III, and $D_1, D_2 < 2$ in Region IV. Point A and B in (b) are tricritical points.

is because $v \leq 0$, indicating an instability of this system. In other words, the system will collapse for sufficiently strong attraction between atoms and molecules.

The 1d ASF can be characterized by the single-particle Green functions of atoms and molecules:

$$\begin{aligned} \langle \Psi_a(\mathbf{x}) \Psi_a^\dagger(0) \rangle &= A_1 \left(\frac{a_0}{r} \right)^{\alpha_1} + \dots, \\ \langle \Psi_m(\mathbf{x}) \Psi_m^\dagger(0) \rangle &= A_2 \left(\frac{a_0}{r} \right)^{\alpha_2} + \dots, \end{aligned} \quad (7)$$

where $\mathbf{x} = (\tau, x)$, $r = \sqrt{(v_+ \tau)^2 + x^2}$, $\Psi_a(x) = a_i / \sqrt{a_0}$, $\Psi_m(x) = b_i / \sqrt{a_0}$, $A_{1,2}$ are nonuniversal constants, and

$$\alpha_2 = 4\alpha_1 = \frac{1}{2K}. \quad (8)$$

We would like to stress that this exact relation between α_1 and α_2 [Eq. (8)] results from the fact that the phase fluctuations of atoms and molecules are locked by the Josephson coupling, and is a characteristic of the 1d ASF. On the other hand, the density correlation functions of atoms and molecules at nonzero momenta decay exponentially due to the presence of the gap Δ_0 . Equation (8) implies that both the atomic and molecular sectors exhibit the behavior of a 1d SF as long as $K > 1/4$. For $1 \leq K_{a/m} < +\infty$, this condition is satisfied, which can be verified from Eq. (6). It gives further support to our claim that this phase is a 1d analog of the ASF in 3d.

In the regime where the g_1 term is irrelevant, the excitation spectrum consists of two branches of gapless excitations with linear dispersion relations. This phase, which has been thoroughly discussed in Ref. 6, will be referred to as the 2-LLs. We just mention that in the present situation this phase can exist only by including sufficiently strong repulsions between atoms or molecules. This is because $D_1 < 2$ for on-site interactions only.

Commensurate filling When δ is close to zero, the g_3 term must be retained. For simplicity, we shall consider $\delta = 0$. Let us first define the scaling dimension of the g_3 term:

$$D_2 = K_a + K_m. \quad (9)$$

The g_1 (g_3) term will be relevant if $D_1 < 2$ ($D_2 < 2$). Accordingly, a tree-level scaling analysis suggests the phase diagram in the K_a - K_m space as shown in Fig. 2 (a). $D_1 < 2$ and $D_2 > 2$ corresponds to the 1d ASF, $D_1 > 2$ and $D_2 < 2$ will be referred to as the I-CDW, and $D_1, D_2 > 2$ corresponds to the 2-LL. However, a competition between the two relevant operators, the g_1 and g_3 terms, occur when $D_1, D_2 < 2$. To determine whether Region IV in Fig. 2 (a) corresponds to a new phase or not, we employ the one-loop RG analysis.

By integrating out the fast modes, the one-loop RG equations are given by

$$\frac{dK_a(l)}{dl} = 2 [\lambda_1^2(l) - K_a^2(l) \lambda_3^2(l)], \quad (10)$$

$$\frac{dK_m(l)}{dl} = \frac{1}{2} [\lambda_1^2(l) - 4K_m^2(l) \lambda_3^2(l)], \quad (11)$$

$$\begin{aligned} \frac{d\lambda_1(l)}{dl} &= \left[2 - \frac{1}{K_a(l)} - \frac{1}{4K_m(l)} \right] \lambda_1(l) \\ &\quad - \frac{\lambda_1(l) \lambda_4(l)}{K_a(l) K_m(l)}, \end{aligned} \quad (12)$$

$$\frac{d\lambda_2(l)}{dl} = \frac{\lambda_1^2(l)}{K_a(l) K_m(l)} - 2\lambda_3^2(l), \quad (13)$$

$$\begin{aligned} \frac{d\lambda_3(l)}{dl} &= [2 - K_a(l) - K_m(l)] \lambda_3(l) \\ &\quad - 2K_a(l) K_m(l) \lambda_2(l) \lambda_3(l), \end{aligned} \quad (14)$$

$$\frac{d\lambda_4(l)}{dl} = 2K_a(l) K_m(l) \lambda_3^2(l) - \lambda_1^2(l), \quad (15)$$

with the initial values: $K_a(0) = K_a$, $K_m(0) = K_m$, $\lambda_{1(3)}(0) = \pi g_{1(3)} / (2v_0)$, $\lambda_2(0) = g_2 / (2v_0)$, and $\lambda_4(0) = 0$. By solving Eqs. (10) — (15), one may find three kinds of behaviors of the RG flow of $\lambda_1(l)$ and $\lambda_3(l)$: (i) $\lambda_1(l)$ flows to strong coupling while $\lambda_3(l)$ flows to zero. This is the 1d ASF. (ii) $\lambda_3(l)$ flows to strong coupling while $\lambda_1(l)$ flows to zero. This is the I-CDW. (iii) Both $\lambda_1(l)$ and $\lambda_3(l)$ flow to zero. This is the 2-LL. Thus, Region IV in Fig. 2 (a) shrinks to a transition line between the 1d ASF and I-CDW in the K_a - K_m space as shown in Fig. 2 (b). Both the phase transition between the 1d ASF and 2 LL and that between the I-CDW and 2-LL belong to the KT type. The transition between the 1d ASF and I-CDW is of second order. Further, all these transition lines coincide at two tricritical points, the point A and B in Fig. 2 (b). The very reason why there is no way for both $\lambda_1(l)$ and $\lambda_3(l)$ flowing to strong coupling simultaneously is that the operators $\cos \sqrt{\pi}(\theta_m - 2\theta_a)$ and $\cos \sqrt{4\pi}(\phi_a - \phi_m)$ are exclusive to one another, that is, the field configurations which minimize one perturbation term do not minimize the other. The interplay between these two competing relevant operators then produces a novel quantum phase transition.

In the I-CDW, it is convenient to define new bosonic fields: $\tilde{\phi}_+ = \frac{1}{2}(\phi_a + \phi_m)$, $\tilde{\phi}_- = \phi_a - \phi_m$, $\tilde{\theta}_+ = \theta_a + \theta_m$, and $\tilde{\theta}_- = \frac{1}{2}(\theta_a - \theta_m)$. The $\tilde{\phi}_+$ and $\tilde{\phi}_-$ fields describe the in-phase and out-of-phase density fluctuations, respectively.

Due to the relevant perturbation $\cos\sqrt{4\pi}\tilde{\phi}_-$, the value of $\langle\tilde{\phi}_-\rangle$ is pinned and a gap is opened for the $\tilde{\phi}_-$ sector, while the $\tilde{\phi}_+$ sector is still gapless. On account of this, both the single-particle Green functions of atoms and molecules decay exponentially. On the other hand, the $2\pi\rho_{a/b}$ parts of the density fluctuations for atoms and molecules are enhanced:

$$\langle\rho_{a(b)}(x)\rho_{a(b)}(0)\rangle|_{2\pi\rho_{a(b)}}\sim\left(\frac{a_0}{|x|}\right)^\gamma, \quad (16)$$

with $\gamma < 2$ for $\pi|g_1|/(2v_0), |g_2|/(2v_0) \ll 1$. Here $\rho_{a/b}(x) = n_{a/b,i}/a_0$.

To understand the nature of the ground state of the I-CDW, a simple picture can be obtained in the limit of strong atom-molecule interactions (g_3), where the potential energy (the g_3 term) dominates over quantum fluctuations. In this case, atoms and molecules form a regular lattice (Wigner crystal of hard-core bosons). For $g_3 > 0$, the energy of the repulsion between atoms and molecules

$$g_3 \cos(\sqrt{4\pi}\phi_-) = -g_3 \cos[\sqrt{4\pi}(\phi_a - \phi_m + \sqrt{\pi}/2)],$$

is minimized by a relative phase shift of $\sqrt{\pi}/2$ between atoms and molecules, which corresponds to a shift of the atom (or molecule) lattice by half-a-period.[9] Thus, the I-CDW respects the symmetry of translation by one site, $a_i \rightarrow a_{i+1}$ and $b_i \rightarrow b_{i+1}$, but spontaneously breaks the reflection symmetry about the origin, $a_i \rightarrow a_{-i}$ and $b_i \rightarrow b_{-i}$ (or $\phi_{a/m} \rightarrow -\phi_{a/m}$ and $\theta_{a/m} \rightarrow \theta_{a/m}$).

Experimental signatures We suggest a few methods to detect the above results experimentally. The most important distinction between the 2-LL and the 1d ASF is the behaviors of the density correlators. For the 2-LLs, the $2\pi\rho_{a(b)}$ part of the density correlators of atoms (molecules) exhibits power-law decay. However, the corresponding sector decays exponentially for the 1d ASF. Therefore, a threshold behavior will be observed at the momentum $k = 2\pi\rho_{a(b)}$ in a time-of-flight measurement for atoms (molecules) in the 1d ASF, while such a behavior does not exist in the 2-LL. The other unique feature of the ASF is that the ground state is a coherent state formed by hybridizing the atoms and molecules. Therefore, by suddenly changing the detuning, some kind of Rabi oscillation will be observed between the atomic and molecular condensates. Such a change of detuning can be achieved by applying magnetic field pulses to the ASF.[10] Although there is no true condensate in 1d due to strong phase fluctuations, a similar oscillation can also be observed between the densities of atoms and molecules, which is strongly damped by the phase fluctuations. Further support to the 1d ASF is the examination of Eq. (8), which can be achieved by a measurement of single-particle Green functions of atoms and molecules through the absorption line shape.

Discussions Finally, two points should be addressed here. First of all, when $a_0|\delta|$ is close to some rational




FIG. 3: Phase diagram for $a_0|\delta|$ being some rational number.

number, one must replace the g_3 term in Eq. (2) by the $\tilde{g}_3$ term [Eq. (3)], with the scaling dimension D_2l . A tree-level scaling analysis suggests the phase diagram in the K_a - K_m space as shown in Fig. 3. We note that there is no direct transition between the 1d ASF and I-CDW. Moreover, the I-CDW can be reached only with very small values of $K_{a/m}$. Next, a RG study shows that small velocity anisotropy, i.e. $v_a \neq v_m$, just shifts the phase boundary, and does not affect our conclusions. This is because the OPE's between the additional operators arising from velocity anisotropy and the already existing operators do not generate themselves.

When this work was finished, we were aware of two recent papers dealing with a related system — a Feshbach-resonant atomic Fermi gas in 1d.[11] Their results about the charge sector are similar to ours for the 1d ASF.

The work of Y.-W. Lee is supported by the National Science Council of Taiwan under grant NSC 93-2112-M-029-007. The work of Y.L. Lee is supported by the National Science Council of Taiwan under grant NSC 93-2112-M-018-009.

* Electronic address: ywlee@thu.edu.tw

† Electronic address: yllee@cc.ncue.edu.tw

- [1] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, E.A. Cornell, Science **269**, 198 (1995); K.B. Davis, M. -O. Mewes, M.R. Andrews, N.J. van Druten, D.S. Durfee, D.M. Kurn, and W. Ketterle, Phys. Rev. Lett. **75**, 3969 (1995).
- [2] S. Jochim, M. Bartenstein, A. Altmeyer, G. Hendl, S. Riedl, C. Chin, J. Hecker Denschlag, and R. Grimm, Science **302**, 2101 (2003); M. Greiner, C.A. Regal, and D.S. Jin, Nature **426**, 537 (2003); M.W. Zwierlein, C.A. Stan, C.H. Schunck, S.M.F. Raupach, S. Gupta, Z. Hadzibabic, and W. Ketterle, Phys. Rev. Lett. **91**, 250401 (2003).
- [3] Y. Ohashi and A. Griffin, Phys. Rev. Lett. **89**, 130402 (2002).
- [4] L. Radzihovsky, J. Park, and P.B. Weichman, Phys. Rev. Lett. **92**, 160402 (2004); M.W.J. Romans, R.A. Duine, S. Sachdev, and H.T.C. Stoof, *ibid.* **93**, 020405 (2004); Y.W. Lee and Y.L. Lee, Phys. Rev. B **70**, 224506 (2004).
- [5] K.K. Das, Phys. Rev. Lett. **90**, 170403 (2003).

- [6] M.A. Cazalilla and A.F. Ho, Phys. Rev. Lett. **91**, 150403 (2003); L. Mathey, D.-W. Wang, W. Hofstetter, M.D. Lukin, and E. Demler, *ibid.* **93**, 120404 (2004).
- [7] We emphasize that most of our results can also be applied to the case where the Feshbach-resonant atomic gas is put in a very elongated trap, such as the toroidal one.
- [8] For a review, see, for example, T. Giamarchi, *Quantum Physics in One Dimension*, (Oxford University Press, Oxford, 2004).
- [9] A similar situation also occurs in quantum wires with two nearly equivalent subbands. See, for example, O.A. Starykh, D.L. Maslov, W. Häusler, and L.I. Glazman in: *Low-Dimensional Systems: Interaction and Transport Properties*, (Springer-Verlag, Berlin, 2000).
- [10] S.J.J.M.F. Kokkelmans and M.J. Holland, Phys. Rev. Lett. **89**, 180401 (2002).
- [11] D.E. Sheehy and L. Radzihovsky, cond-mat/0505681; R. Citro and E. Orignac, cond-mat/0505706.